

Assignment 2

(28 February, 2026)

Submission Deadline: 21 March, 2026

Course: Algebraic Topology II (KSM4E02)

Instructor: Aritra Bhowmick

Q1. (Internal Hom) Given two chain complexes $C_\bullet, D_\bullet$, their *internal hom* is defined as

$$[C, D]_n = \prod_{k \in \mathbb{Z}} \text{hom}(X_k, Y_{k+n}).$$

Define $d_n : [C, D]_n \rightarrow [C, D]_{n-1}$ via the formula

$$d_n(h_k) = (\partial_{k+n}^D \circ h_k - (-1)^n h_{k-1} \circ \partial_k^C)$$

Verify the following.

- $[C, D]_\bullet = ([C, D]_n, d_n)$ is a chain complex.
- The n -cycles of $[C, D]_\bullet$ are precisely the degree- n chain maps $C_\bullet \rightarrow D_\bullet$.
- The 0^{th} homology of $[C, D]_\bullet$ consists of the chain homotopy classes of chain maps $C_\bullet \rightarrow D_\bullet$.

Note: Why *internal* hom?! Consider the category Ab . The hom-set $\text{hom}_{\text{Ab}}(X, Y)$ itself is naturally an Abelian group, and thus it is already *internal* to the category. Similarly, in Top , we can use the compact-open topology to make $\text{hom}_{\text{Top}}(X, Y)$ into a topological space. For Ch , we could treat $\text{hom}_{\text{Ch}}(C_\bullet, D_\bullet)$ as a chain complex concentrated at degree 0. The internal hom functor $[C_\bullet, _]$ is right adjoint to the tensor functor $_ \otimes C_\bullet$; this is essentially the categorical definition of internal hom. We shall later see the definition of the tensor product of chain complexes.

3 + 3 + 4 = 10

Q2. (Good Pair) A pair (X, A) is called a *good pair* if $A \subset X$ is closed, and there exists an open neighborhood $A \subset U \subset X$ such that U deformation retract onto A . Given a good pair (X, A) , consider the diagram

$$\begin{array}{ccccc} H_n(X, A) & \xrightarrow{\text{blue}} & H_n(X, U) & \xleftarrow{\text{green}} & H_n(X \setminus A, U \setminus A) \\ \downarrow q_\star & & \downarrow q_\star & & \downarrow q_\star \\ H_n(X/A, A/A) & \xrightarrow{\text{blue}} & H_n(X/A, U/A) & \xleftarrow{\text{green}} & H_n(X/A \setminus A/A, U/A \setminus A/A) \end{array}$$

where q is the quotient map that identifies A to a point. Verify that the diagram is commutative. Show that $q_\star : H_n(X, A) \rightarrow H_n(X/A, A/A)$ is an isomorphism.

Hint : The *green* arrows are isomorphism by excision. For the *blue* arrow in the top row, note that $H_n(U, A) = 0$, and then use the long exact sequence for the triple (X, U, A) ; similar argument works for the bottom row as well. The right-most *purple* vertical arrow is induced by a homeomorphism

3 + 7 = 10


Q3. (Homology Long Exact Sequence of Cofibration) Let $f : A \rightarrow X$ be a cofibration with the mapping cone C_f . Show that there exists a long exact sequence (in any homology theory)

$$\cdots \rightarrow \tilde{H}_{n+1}(C_f) \rightarrow \tilde{H}_n(A) \rightarrow \tilde{H}_n(X) \rightarrow \tilde{H}_n(C_f) \rightarrow H_{n-1}(A) \rightarrow \cdots$$

Hint : Replace $f : A \rightarrow X$ by the mapping cylinder construction

10

Q4. (Lens Space) Construct a three-dimensional CW complex as follows. Take n -many 3-simplices $T_1, \dots, T_n$. First, join them cyclically in the following way:



Next, identify the bottom face of T_i with the top face of T_{i+1} , where the indices are taken mod n (so that the bottom face of T_n is identified with the top face of T_1). Denote this space by L_n . Show that the singular homology groups are

$$H_i(L_n) = \begin{cases} \mathbb{Z}, & i = 0 \\ \mathbb{Z}/n\mathbb{Z}, & i = 1 \\ 0, & i = 2 \\ \mathbb{Z}, & i = 3 \\ 0, & i \geq 4. \end{cases}$$

This space is an example of a *lens space*.

10